

Diabetes and Chronic Kidney Disease in Türkiye (DIAKIT): A Cross-Sectional Cohort Study

Mustafa Arici, Kenan Ates, Alaattin Yildiz, Ali R. Odabas, Bulent Tokgoz, Siren Sezer; Halil Zeki Tonbul, Elif Ari Bakir, Memnune Sena Ulu, Ozkan Gungor, Nurhan Seyahi, Sim Kutlay, Bulent Altun

Correspondence to: Mustafa Arici, MD, Prof; E-mail: marici@hacettepe.edu.tr

Table S3. Glomerular filtration rate (GFR) stages according to sex

		Sex			p*
		Male	Female	Total	
GFR Stage, n (%)	Stage 1 (90.0 $\geq$)	245 (45.2)	588 (57.0)	833 (52.9)	<0.001
	Stage 2 (60.0-89.9)	229 (42.3)	332 (32.2)	561 (35.6)	
	Stage 3A (45.0-59.9)	52 (9.6)	74 (7.2)	126 (8.0)	
	Stage 3B (30.0-44.9)	14 (2.6)	30 (2.9)	44 (2.8)	
	Stage 4 (15.0-29.9)	1 (0.2)	4 (0.4)	5 (0.3)	
	Stage 5 (15.0 $\leq$)	1 (0.2)	4 (0.4)	5 (0.3)	
	Total	542 (100.0)	1032 (100.0)	1574 (100.0)	

* Fisher's Exact Test

Table S4. Glomerular Filtration rate (GFR) stages according to age groups

		Age Group						Total
		18-39	40-49	50-59	60-69	70-79	80+	
GFR Stage, n (%)	Stage 1 (90.0 $\geq$)	17 (94.4)	101 (91.0)	313 (75.1)	327 (50.9)	73 (22.1)	2 (3.7)	833 (52.9)
	Stage 2 (60.0-89.9)	1 (5.6)	8 (7.2)	87 (20.9)	249 (38.7)	187 (56.5)	29 (53.7)	561 (35.6)
	Stage 3A (45.0-59.9)	0 (0.0)	2 (1.8)	13 (3.1)	45 (7.0)	52 (15.7)	14 (25.9)	126 (8.0)
	Stage 3B (30.0-44.9)	0 (0.0)	0 (0.0)	2 (0.5)	18 (2.8)	16 (4.8)	8 (14.8)	44 (2.8)
	Stage 4 (15.0-29.9)	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.3)	3 (0.9)	0 (0.0)	5 (0.3)
	Stage 5 (15.0 $\leq$)	0 (0.0)	0 (0.0)	2 (0.5)	2 (0.3)	0 (0.0)	1 (1.9)	5 (0.3)
	Total	18 (100.0)	111 (100.0)	417 (100.0)	643 (100.0)	331 (100.0)	54 (100.0)	1574 (100.0)

Table S5. Glomerular filtration rate (GFR) values of the patients according to sex

		N	Mean±SD Median (Q1-Q3) (Min.-Max.)	p*
GFR	Male	542	84.41±18.17 88.50 (74.00-96.90) (9.40-124.90)	<0.001
	Female	1032	87.59±19.33 92.50 (77.85-100.10) (8.60-134.30)	
	Total	1574	86.50±19.00 90.95 (76.50-99.00) (8.60-134.30)	

*Mann-Whitney U test