

**The incidence, risk factors and maternal and foetal outcomes of uterine rupture
during different birth policy periods: an observational study in China**

Yangwen Zhou¹, Yi Mu², Peiran Chen¹, Yanxia Xie¹, Jun Zhu^{1,†} and Juan Liang^{1,†}

Additional file 1. Number of uterine ruptures by maternal age, parity and number previous caesarean sections over different birth policy periods in 438 hospitals, China.

Maternal characteristics	One-child policy period	Partial two-child policy period	Universal two-child policy period
Number of uterine ruptures (%)			
Maternal age			
<24	242 (14.68)	344 (10.87)	519 (6.39)
24-29	588 (35.68)	1052 (33.25)	2309 (28.43)
30-34	528 (32.04)	1121 (35.43)	3181 (39.17)
≥35	233 (14.14)	553 (17.48)	2045 (25.18)
Missing	57 (3.46)	94 (2.97)	68 (0.84)
Parity			
0	274 (16.63)	266 (8.41)	408 (5.02)
1	1100 (66.75)	2397 (75.76)	6201 (76.35)
2	237 (14.38)	452 (14.29)	1277 (15.72)
≥3	36 (2.18)	49 (1.55)	140 (1.72)
Missing	1 (0.06)	0 (0.00)	96 (1.18)
Previous caesarean sections			
0	411 (24.94)	411 (12.99)	723 (8.90)
1	1090 (66.14)	2434 (76.93)	6273 (77.23)
≥2	139 (8.43)	312 (9.86)	1027 (12.64)
Missing	8 (0.49)	7 (0.22)	99 (1.22)
Total	1648 (100.00)	3164 (100.00)	8122 (100.00)