

Table S2. Percentage of subpopulations defined by the combination of stem cell markers CD44 and CD24 in MCF-7 control cells or treated with 4m, 4bt (dimethylenastron), 4p, 4bc, 4x or monastrol for 24 h. Data represent the mean $\pm$ SEM of 3 independent experiments.

MCF-7		PHENOTYPES (MEAN $\pm$ SEM)			
Group		CD44⁻/CD24⁻	CD44⁻/CD24⁺	CD44⁺/CD24⁻	CD44⁺/CD24⁺
Control		2.14 $\pm$ 0.15	65.21 $\pm$ 9.23	0.38 $\pm$ 0.18	32.26 $\pm$ 9.20
4m (1.0 mM)		1.01 $\pm$ 0.12	55.59 $\pm$ 15.09	0.28 $\pm$ 0.15	43.10 $\pm$ 15.06
4bt (0.8 mM)		3.09 $\pm$ 0.46	34.92 $\pm$ 19.45	0.005 $\pm$ 0.005	61.97 $\pm$ 19.91
4p (0.4 mM)		2.83 $\pm$ 0.59	66.42 $\pm$ 10.57	0.33 $\pm$ 0.20	30.40 $\pm$ 9.78
4bc (1.0 mM)		3.90 $\pm$ 0.070	46.49 $\pm$ 18.49	0.18 $\pm$ 0.07	49.42 $\pm$ 18.49
4x (0.8 mM)		2.45 $\pm$ 0.52	61.35 $\pm$ 20.83	1.065 $\pm$ 0.97	35.11 $\pm$ 19.33
Monastrol (1.0 mM)		1.64 $\pm$ 0.43	68.56 $\pm$ 5.93	0.2400 $\pm$ 0.20	29.53 $\pm$ 5.30