

Supplementary Table S4. Characteristics of patients with endometriosis and patients with adenomyosis.

Patient Characteristics		Adenomyosis only (n=59)	Endometriosis (n=117)	Alpha
Age (years) Mean $\pm$ SD		53.5 $\pm$ 9.7	46.0 $\pm$ 12.0	<.001
Premenopausal		29 (49.2)	79 (67.5)	.018
Epithelial Ovarian Cancer		41 (69.5)	70 (59.8)	.210
	Serous HG	25 (42.4)	18 (15.4)	<.001
	Serous LG	3 (5.1)	7 (6.0)	.808
	Endometrioid LG	4 (6.8)	22 (18.8)	.034
	Endometrioid HG	2 (3.4)	5 (4.3)	.777
	Clear cell	2 (3.4)	14 (12.0)	.062
Borderline		11 (18.6)	41 (35.0)	.024
Stage	1	20 (33.9)	75 (64.1)	<.001
	2	4 (6.8)	13 (11.1)	
	3	20 (33.9)	14 (12.0)	
	4	12 (20.3)	5 (4.3)	
ECOG Status	0	37 (62.7)	100 (85.5)	<.001
	1	18 (30.5)	9 (7.7)	
Surgery	Primary	54 (91.5)	115 (98.3)	.030
	Interval	5 (8.5)	2 (1.7)	
Complete Resection		54 (96.4)	110 (94.0)	.536
Grading	G1-G2	7 (11.9)	34 (29.1)	.002
	G3	34 (57.6)	40 (34.2)	
	Gx	18 (30.5)	43 (36.8)	
p53 *		26 (72.2)	20 (32.8)	<.001
BRCA *		4 (20.0)	3 (7.3)	.145

Intergroup comparisons between patients with adenomyosis only and patients with endometriosis were performed using the Pearson χ^2 test for categorical variables and independent t-test for continuous variables. Abbreviations: ECOG = Eastern Cooperative Oncology Group; HG = high-grade; LG = low-grade; SD = standard deviation. * relative percentages.