

Supplemental Table 1: Survival Analysis Assessing Factors Associated with Overall Survival (N=158)

		Unadjusted HR	P-value	Adjusted HR	P-value
Stage	Localized disease	Ref			
	Metastatic	1.79 (1.15, 2.78)	0.01	1.92 (1.23, 3.01)	0.004 *
Gender	Female gender	Ref			
	Male gender	1.71 (1.10,2.66)	0.017	1.84 (1.18, 2.87)	0.007 *
Pathological Grade	Well-diff	Ref			
	Mod/poor diff	1.34 (0.71, 2.55)	0.37		
	Unknown	1.30 (0.71, 2.38)	0.39		
Disease site	Colon cancer	Ref			
	Rectal cancer	1.13 (0.65,1.95)	0.67		
Home location	Rural	Ref			
	Urban	0.97 (0.60,1.56)	0.91		
Traditional medicine use	No traditional medicine	Ref			
	Traditional medicine	1.18 (0.73,1.90)	0.49		
Age	Age<50	Ref			
	Age>=50	0.98 (0.63,1.54)	0.94		

Supplemental Table 2: Survival Analysis Assessing Factors Associated with Disease-Free Survival in patients with localized disease (N=96)

		Unadjusted HR (95% CI)	P-value
Disease site	Colon cancer	Ref	
	Rectal cancer	2.35 (1.01,5.48)	0.047
Gender	Female gender	Ref	
	Male gender	1.19 (0.72,1.97)	0.49
Pathological Grade	Well-diff	Ref	
	Mod/poor diff	1.61 (0.80, 3.23)	0.18
	Unknown	1.35 (0.70, 2.59)	0.37
Home location	Rural	Ref	
	Urban	0.88 (0.50,1.55)	0.66
Traditional medicine use	No traditional medicine	Ref	
	Traditional medicine	0.79 (0.46,1.35)	0.40
Age	Age<50	Ref	
	Age>=50	1.40 (0.83,2.39)	0.21

Supplemental Figure 1: Cohort Ascertainment

