




(A) Depressive episode




(B) Mixed episode



(C) (Hypo)manic episode



(D) Total sample



SI: Intensity of suicidal ideation
SA: Occurrence of suicide attempts (in the past month)
DEP: Score of the depressive subscale of CUDOS-M
MAN: Score of the (hypo)manic subscale of CUDOS-M
EPI: Number of mood episodes

NSSI: Frequency of non-suicidal self-injury (in the past month)
ADD 1: The self-injury occurs more often than intended.
ADD 2: The severity in which self-injury occurs has increased.
ADD 3: To self-injure more frequently or with greater intensity to produce the same effect.
ADD 4: The self-injury consumes a significant amount of time.
ADD 5: Despite a desire to control self-injury, one is unable to do so.
ADD 6: To continue self-injury despite recognizing that it is harmful physically and/or emotionally.
ADD 7: Important social, family, academic, or recreational activities are given up or reduced because of self-injury.

Supplemental Fig. S2 Network structures of suicidality and NSSI based on the Akaike Information Criterion

(A) Depressive episode; (B) Mixed episode; (C) (Hypo)manic episode; (D) Total sample.

CUDOS-M, the Clinically Useful Depression Outcome Scale supplemented with questions for the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5) mixed subtype.

For the network structures, edge thickness represents the degree of association. Green edges indicate positive association, and red edges indicate negative association.