

BMCAdditionalfile1 Prevalence of Limb Pain by Sociodemographic Variables in Icelandic Schoolchildren

	Frequency of Extremity Pain (%)						
	Almost Daily	More than once a week	Once a week	At least once a month	Rarely or never	Chi Squared (df)	p- value
Sex^a							
Boys (n=5304)	4.4	6.3	10.6	18.6	60.1	14.2(4)	.007
Girls (n=5254)	5.3	7.8	10.2	18.3	58.4		
Grade^b							
6 (n=3404)	4.3	6.4	8.8	15.9	64.5	70.3(8)	<.001
8 (n=3676)	5.0	7.0	11.9	20.4	55.7		
10 (n=3439)	5.5	7.8	10.4	18.7	57.6		
Family Structure^c							
Both parents (n=7241)	4.4	6.8	10.6	18.1	60.2	62.5(12)	<.001
Stepparent (n=1408)	5.8	7.1	10.8	19.4	56.9		
Single parent (n=1565)	5.1	8.6	9.3	18.6	58.4		
Other arrangement (n=183)	15.3	5.5	11.5	13.7	54.1		
Family Affluence^d							
Low (n=2640)	5.8	6.8	10.2	17.5	59.7	10.1(8)	ns
Medium (n=4743)	4.4	7.3	10.8	18.5	59.0		
High							

(n=2420)	4.5	6.9	10.3	19.0	59.2
----------	-----	-----	------	------	------

^a = 2% missing data, ^b = 2.3% missing data, ^c = 3.5% missing data, ^d = 9.0% missing data