

Additional file 5. Change of IIEF and EDITS total scores from baseline to Month 6 during treatment with tadalafil OaD - subgroup analysis (N=778; unadjusted data)

Mean (SD) score change from baseline to Month 6						
Variable	IIEF domain scores					EDITS total score ^a n=183
	Erectile function n=776	Intercourse satisfaction n=776	Orgasmic function n=776	Overall satisfaction n=776	Sexual desire n=776	
By age						
Age ≤65 years	8.6 (7.9)	3.7 (4.1)	1.6 (2.9)	2.9 (2.9)	1.0 (2.0)	17.9 (23.3)
Age >65 years	8.0 (7.3)	3.3 (3.8)	1.6 (3.4)	2.8 (2.9)	0.8 (2.2)	11.5 (17.5)
By PDE5-I treatment						
PDE5-I naïve	9.1 (7.7)	3.9 (3.9)	1.8 (3.0)	3.1 (2.8)	1.0 (2.0)	4.5 (17.7)
PDE5-I pretreated	7.4 (7.9)	3.1 (4.2)	1.3 (3.0)	2.5 (3.0)	0.9 (2.1)	16.6 (22.1)
By disease severity						
Mild ED	6.2 (6.9)	2.8 (4.1)	1.2 (2.9)	2.2 (2.8)	0.6 (1.9)	10.9 (19.5)
Moderate ED	8.6 (7.6)	3.7 (3.6)	1.5 (2.9)	2.9 (2.8)	1.1 (2.0)	15.2 (21.1)
Severe ED	10.2 (8.6)	4.2 (4.8)	2.5 (3.3)	3.5 (3.0)	1.0 (2.2)	20.3 (24.4)
By presence of BPH						
BPH present	5.7 (6.7)	2.5 (3.2)	0.8 (3.1)	2.5 (2.8)	0.0 (2.2)	14.2 (18.5)
BPH absent	8.7 (7.9)	3.7 (4.1)	1.7 (3.0)	2.9 (2.9)	1.0 (2.0)	16.5 (22.4)
By presence of diabetes						
Diabetes present	9.8 (7.6)	4.0 (4.2)	2.2 (3.4)	3.1 (2.9)	0.9 (1.9)	21.6 (22.9)
Diabetes absent	8.2 (7.8)	3.6 (4.0)	1.5 (3.0)	2.8 (2.9)	0.9 (2.1)	15.1 (21.8)
By presence of CVD						
CVD present	8.2 (8.0)	3.3 (4.0)	1.6 (3.4)	2.6 (3.0)	1.0 (2.1)	17.8 (24.3)
CVD absent	8.6 (7.7)	3.8 (4.1)	1.7 (2.8)	3.0 (2.8)	0.9 (2.0)	15.3 (20.6)
By presence of hypertension						
Hypertension present	8.3 (8.0)	3.3 (4.0)	1.7 (3.4)	2.6 (3.0)	1.0 (2.1)	18.3 (24.2)
Hypertension absent	8.6 (7.7)	3.8 (4.0)	1.6 (2.8)	3.0 (2.8)	0.9 (2.0)	15.0 (20.6)
By presence of dyslipidemia						
Dyslipidemia present	8.3 (8.3)	3.4 (4.3)	1.5 (3.3)	2.7 (2.9)	0.8 (2.1)	13.6 (25.3)
Dyslipidemia absent	8.5 (7.7)	3.7 (4.0)	1.7 (3.0)	2.9 (2.9)	1.0 (2.0)	17.0 (21.2)

^a EDITS scores at baseline were collected only in patients pretreated with PDE5-I

BPH, benign prostatic hyperplasia; CVD, cardiovascular disease; ED, Erectile Dysfunction; EDITS, Erectile Dysfunction Inventory of Treatment Satisfaction; IIEF, International Index of Erectile Function; N, number of patients; n, number of patients with data at baseline; OaD, once a day; PDE5-I, phosphodiesterase type 5 inhibitor; SD, standard deviation