

Impacted Mandibular Third Molar Difficulty Assessment Questionnaire

Patient Information

Patient Name / Surname: _____

File Number: _____

Date: _____

Tooth Location: Right () Left ()

SYSTEMIC ASSESSMENT

1. Presence of systemic disease: Yes (1) No (0)
2. Presence of diagnosed psychological condition: Yes (1) No (0)
3. History of psychiatric medication use: Yes (1) No (0)

RADIOGRAPHIC ASSESSMENT

1. Relationship with the mandibular canal:

- Distance $\geq$ 2 mm (1)
- Distance < 2 mm (2)
- Root in contact with the canal (3)
- Canal located between the roots (4)

2. Number of roots:

- Single root (1)
- Two roots (2)
- More than two roots (3)

3. Root morphology:

- Severe root curvature (1)
- Thick septum (1)

4. Angulation and tooth position:

- Mesioangular (1)
- Horizontal (2)

- Vertical (3)
- Distoangular (4)
- Buccolingual (5)
- Inverted (6)

5. Ramus relationship (based on space):

- Class I ($M3 > \text{mesiodistal width}$) (1)
- Class II ($M3 = \text{mesiodistal width}$) (2)
- Class III ($M3 < \text{mesiodistal width}$) (3)

6. Depth relative to second molar:

- Crown to crown (1)
- Crown to cementoenamel junction (2)
- Crown to root (3)

7. Presence of adjacent tooth: Yes (0) No (1)

8. Presence of pathological findings: Yes (1) No (0)

CLINICAL ASSESSMENT

1. Age:

- 0–29 (1)
- 30–39 (2)
- >40 (3)

2. Maximum interincisal opening:

- 30–39 mm (3)
- 40–49 mm (2)
- >50 mm (1)

3. Presence of pericoronitis: Yes (1) No (0)

4. Eruption status:

- Partially erupted (1)

- Fully impacted (2)

5. Body Mass Index (BMI):

- Normal weight (0)

- Overweight (1)

POSTOPERATIVE ASSESSMENT

Operation duration: _____

Complications: Infection () Trismus () Root displacement ()