

Demography

Please complete the survey below.

Thank you!

3% Progress

Phase 3 Delphi Survey

Title: Identifying Quality Indicators for End-of-Life Cancer Care in Malaysia: a modified Delphi study

Dear participant,

We kindly invite you as an expert panel to participate in our Phase 3 survey, to help us by giving your opinion on the feasibility of extracting those quality indicators from patient medical records based on the numerator and denominator given.

The goal of this study is to develop an expert-panel endorsed list of quality indicators (QIs) for end-of-life care in adult cancer patients (≥ 18 years of age) that is applicable in Malaysia settings. Quality indicators can indicate potential problems (such as appropriate or inappropriate treatment) or good quality care. End-of-life care refers to all aspects of care that is provided during the period when death is about to happen. There is no general agreement regarding the end-of-life time frame: definitions range from care in the last hours or months or last year of life, to care from the time of terminal diagnosis until death.

*The link below contains detailed information on the study flow chart.

[Attachment: "Study Overview3.pdf"]

Your Particulars:

We greatly appreciate it if you could kindly leave your contact details which will be used for tracking purposes.

It will be kept strictly confidential and we will not contact you about anything other than the study.

Email address:

Contact number:

((e.g, 0125163137, without "-"))

Phase 2, Round 2 Results are here!!!

Feel free to view the analysis by clicking the attachment link below!

Summary results of Phase 1 & 2 surveys: - Phase 1: 41 out of a total of 64 quality indicators provided had reached consensus & 2 new indicators moved to next phase.

- Phase 2 (Round 1): 38 out of 43 quality indicators from phase 1 had reached consensus and moved to phase 3. The remaining 5 quality indicators that did not reached consensus were presented again in the subsequent round.

- Phase 2 (Round 2): 2 out of 5 quality indicators from round 1 of phase 2 had reached consensus. A total of 40 quality indicators that have reached consensus from phase 2 moved to this phase 3 survey .

[Attachment: "Phase 2 (round 2) results.pdf"]

Phase 3: Pain

Please complete the survey below.

Thank you!

12% Progress

Instructions:

- As you complete the survey, we ask that you keep in mind that the patient population being considered is only adult patients with cancer.
- Please answer each question based on your knowledge and personal experiences in end-of-life cancer care services you most often interact with in Malaysia.
- Please provide an answer to each statement and elaborate on your answer by putting down your comments (if necessary). This will help us to better understand the answer you provide.
- Please use the 9-point Likert scale to rate your answer. A score of "1" indicates that the quality indicator of end-of-life cancer care is not at all feasible and a score of "9" indicates highly feasible to be extracted from medical records. Please consider your rating based on the information in the boxes (i.e., the numerator and denominator) whether this information would be available in medical records/databases rather than if the quality indicators can be achieved in our healthcare system.

Please complete the survey below:

PAIN

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients with documented pain assessment in the last 1 month before death

Numerator: No. of patients with documented pain assessment in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients prescribed with pain management (pharmacological and/or non-pharmacological approach) in the last 1 month before death

Numerator: No. of patients prescribed with pain management in the last 1 month before death

Denominator: No. of patients who died from cancer and with documented pain

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who received opioids in the last 1 month before death

Numerator: No. of patients who received opioids in the last 1 month before death

Denominator: No. of patients who died from cancer and with moderate to severe pain

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who were treated with opioids and without bowel obstruction received concurrent stool softener in the last 1 month before death

Numerator: No. of patients who were treated with opioids received concurrent prophylaxis of Opiate-Induced Constipation (OIC) in the last 1 month before death

Denominator: No. of patients who died from cancer and on opioids and without contraindications to bowel regimen

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who were prescribed neuropathic medications for their nerve pain in the last 6 months before death

Numerator: No. of patients prescribed with medications for neuropathic pain in the last 6 months before death

Denominator: No. of patients who died from cancer and with documented neuropathic pain

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: DYSPNOEA

Please complete the survey below.

Thank you!

24% Progress

DYSPNOEA

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients who had documented dyspnoea assessment in the last 1 month before death

Numerator: No. of patients with documented dyspnoea assessment in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients received treatment for breathlessness (pharmacological and/or non-pharmacological approach) in the last 2 weeks before death

Numerator: No. of patients received treatment for dyspnoea in the last 2 weeks before death

Denominator: No. of patients who died from cancer and with documented dyspnoea

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: OTHER PHYSICAL SYMPTOMS

Please complete the survey below.

Thank you!

32% Progress

OTHER PHYSICAL SYMPTOMS

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients with documented assessment of nausea or vomiting in the last 1 month before death

Numerator: No. of patients with documented assessment of nausea or vomiting in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had documented oral care in the last 1 month before death

Numerator: No. of patients who had documented oral care in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients with documented assessment of delirium and agitation in the last 2 weeks before death

Numerator: No. of patients with documented assessment of delirium and agitation in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients with delirium and agitation received treatment (pharmacological and/or non-pharmacological approach) in the last 2 weeks before death

Numerator: No. of patients with delirium and agitation received treatment in the last 2 weeks before death

Denominator: No. of patients who died from cancer and with documented delirium/agitation

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients with distressing symptoms treated with rescue medications in the last 2 weeks before death

Numerator: No. of patients treated with rescue medications in the last 2 weeks before death

Denominator: No. of patients who died from cancer and with documented distressing symptoms

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: PSYCHOLOGICAL AND SOCIAL ASPECTS OF CARE

Please complete the survey below.

Thank you!

40% Progress

PSYCHOLOGICAL AND SOCIAL ASPECT OF CARE

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients with documented psychological assessment in the last 6 months before death

Numerator: No. of patients with documented psychological assessment in the last 6 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients with depressive symptoms had documented assessment of depression in the last 6 months before death

Numerator: No. of patients with documented assessment of depression in the last 6 months before death

Denominator: No. of patients who died from cancer and with depressive symptoms

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients with depression received treatment (pharmacological and/or non-pharmacological approach) in the last 6 months before death

Numerator: No. of patients with received treatment for depression in the last 6 months before death

Denominator: No. of patients who died from cancer and diagnosed with depression

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients with documented social needs assessment (e.g. family support, transportations, conditions related to living arrangement, financial status and etc) in the last 6 months before death

Numerator: No. of patients with documented social needs assessment in the last 6 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients offered social support and documented in the last 3 months before death

Numerator: No. of patients offered social support and documented in the last 3 months before death

Denominator: No. of patients who died from cancer and with documented social needs

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: SPIRITUAL CARE

Please complete the survey below.

Thank you!

50% Progress

SPIRITUAL CARE

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients had spiritual needs assessment in the last 6 months before death

Numerator: No. of patients with documented spiritual needs assessment in the last 6 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients offered spiritual support and documented in the last 6 months before death

Numerator: No. of patients offered spiritual support and documented in the last 6 months before death

Denominator: No. of patients who died from cancer and with documented spiritual needs

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: TREATMENT MODALITIES

Please complete the survey below.

Thank you!

55% Progress

TREATMENT MODALITIES

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients received antiemetics for chemotherapy in the last 6 months before death

Numerator: No. of patients who received antiemetics for chemotherapy in the last 6 months before death

Denominator: No. of patients who died from cancer and received chemotherapy

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: HOSPITAL & COMMUNITY PALLIATIVE CARE SERVICES

Please complete the survey below.

Thank you!

60% Progress

HOSPITAL & COMMUNITY PALLIATIVE CARE SERVICES

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients received first referral to community palliative care services in the last 6 months before death

Numerator: No. of patients who received first referral to community palliative care services in the last 6 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients received first referral to hospital palliative care unit in the last 6 months before death

Numerator: No. of patients who received first referral to hospital palliative care unit in the last 6 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who received home care visits (scheduled) in the last 3 months before death

Numerator: No. of patients who received home care visits in the last 3 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had one or more doctor house calls (unscheduled) in the last 2 weeks before death

Numerator: No. of patients who had one or more doctor house calls (unscheduled) in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients' families who received home visits or phone calls within 2 weeks after patients' death**Numerator: No. of patients' families who received home visits or phone calls within 2 weeks after patients' death****Denominator: No. of patients who died from cancer and with documented bereavement needs for their family/caregivers**

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of doctors trained in basic palliative care (i.e., to provide basic symptom assessment and relief for patients with life-limiting conditions; equipped with knowledge on when/how to refer patients for specialised care)**Numerator:****No. of doctors trained in basic palliative care****Denominator: No. of doctors in the hospital/institute/organisation**

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of nurses with at least advanced diploma in palliative care**Numerator: No. of nurses with at least advanced diploma in palliative care****Denominator: No. of nurses involved in palliative care services**

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: ADVANCED CARE PLANNING (ACP)

Please complete the survey below.

Thank you!

65% Progress

ADVANCED CARE PLANNING (ACP)

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients who have documented key decision-maker in the last 4 weeks before death

Numerator: No. of patients with documented key decision-maker in the last 4 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who have documented resuscitation preferences in the last 2 weeks before death

Numerator: No. patients with documented resuscitation preferences in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3 CONTINUITY & COORDINATION OF CARE

Please complete the survey below.

Thank you!

75% Progress

CONTINUITY & COORDINATION OF CARE

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients who had documented multidisciplinary medical consultation in the last 3 months before death

Numerator: No. of patients with documented multidisciplinary medical consultation in the last 3 months before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had documented explanations of their medical conditions to them in the last 1 month before death

Numerator: No. of patients with documented explanations of their medical conditions to them in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had documented explanations of their impending death to them in the last 1 month before death

Numerator: No. of patients who had documented explanations of their impending death to them in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients whose family had documented explanation of patient's medical condition in the last 1 month before death

Numerator: No. of patients whose family had documented explanation of patient's medical condition in the last 1 month before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients whose family had documented explanation of patient's impending death in the last 2 weeks before death

Numerator: No. of patients whose family had documented explanation of patient's impending death in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: PLACE OF CARE / DEATH

Please complete the survey below.

Thank you!

87% Progress

PLACE OF CARE/DEATH

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients who had documented discharge plans in the last 2 weeks before death

Numerator: No. of patients with documented discharge plans in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had their preferred place of care documented in the last 2 weeks before death

Numerator: No. of patients with their preferred place of care documented in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had their families' preferred place of care for the patient documented in the last 2 weeks before death

Numerator: No. of patients with their families' preferred place of care documented in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who had their preferred place of death documented in the last 2 weeks before death

Numerator: No. of patients with their preferred place of death documented in the last 2 weeks before death

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Percentage of patients who died at home

Numerator: No. of patients who died at home

Denominator: No. of patients who died from cancer

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Phase 3: MEDICATION

Please complete the survey below.

Thank you!

93% Progress

MEDICATION

How feasible is it for this indicator to be extracted from the medical records?

FEASIBILITY

(1= not at all feasible; 9 = highly feasible)

Percentage of patients receiving medication for respiratory secretion in the last 2 weeks before death

Numerator: No. of patients prescribed with medication for respiratory secretion in the last 2 weeks before death

Denominator: No. of patients who died from cancer with documented respiratory secretion

Feasibility:

((1= not at all feasible; 9 = highly feasible))

☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5 ☐ 6 ☐ 7 ☐ 8 ☐ 9

If you have any other comments regarding this quality indicator, please kindly leave them here.

Thank you for your time & patience in doing this survey.

We hope to get back to you in 4 weeks to participate in our last round of survey.
THANK YOU!