

Comparison of the 24 h and 48 h effects of extracts of *Olax subscorpioidea* (OS) and *Uvaria chamae* (UC) on the viability of HeLa cells (n=3)

Conc. ($\mu\text{g/ml}$)	OS		UC	
	24 h	48 h	24 h	48 h
0	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00	100.00 $\pm$ 0.00
5	101.62 $\pm$ 4.80	101.58 $\pm$ 1.00	111.46 $\pm$ 2.77	94.36 $\pm$ 2.14
10	99.28 $\pm$ 1.71	104.61 $\pm$ 2.56	104.24 $\pm$ 3.68	98.61 $\pm$ 4.27
15	91.19 $\pm$ 4.09	98.19 $\pm$ 2.67	113.86 $\pm$ 5.58	102.12 $\pm$ 5.68
25	82.63 $\pm$ 5.37	85.15 $\pm$ 9.81	104.38 $\pm$ 3.91	102.02 $\pm$ 2.42
50	75.10 $\pm$ 2.68	75.25 $\pm$ 12.82	97.57 $\pm$ 2.91	101.20 $\pm$ 5.71
75	73.16 $\pm$ 2.70	71.26 $\pm$ 14.22	93.62 $\pm$ 3.12	96.73 $\pm$ 8.30
100	72.88 $\pm$ 2.74	67.73 $\pm$ 15.31	84.32 $\pm$ 3.77	88.81 $\pm$ 11.73
200	75.46 $\pm$ 2.68	70.41 $\pm$ 14.11	51.76 $\pm$ 11.96	42.69 $\pm$ 26.22
500	19.58 $\pm$ 2.28	20.53 $\pm$ 3.69	10.92 $\pm$ 0.95	7.68 $\pm$ 0.55