

Additional file 1

Theory of prokaryotic pangenome evolution

Itamar Sela, Yuri I. Wolf, and Eugene V. Koonin

Genomes names	ATGC	# genomes	# slow genes	# fast genes	P_{slow}^-	P_{fast}^-
Escherichia coli/Salmonella enterica	ATGC001	20	2929	1471	563	3877
Klebsiella/Enterobacter aerogenes	ATGC002	20	3558	1648	502	7962
Bacillus thuringiensis/cereus/anthracis	ATGC014	20	4057	1497	519	4081
Bacillus subtilis/amyloliquefaciens/velezensis	ATGC015	20	3178	658	195	679
Mycoplasma mycoides/capricolum	ATGC035	17	504	319	12	341
Rickettsia rickettsii/conorii/massiliae	ATGC044	15	768	429	581	20706
Ehrlichia chaffeensis/ruminantium/muris	ATGC045	14	814	62	102	550
Helicobacter pylori/cetorum	ATGC050	20	1232	222	187	812
Corynebacterium glutamicum/deserti/callunae	ATGC069	14	1828	1002	2	625
Pseudomonas putida/montelii/parafulva	ATGC071	20	3452	1636	278	4457
Pseudomonas syringae/savastanoi/cichorii	ATGC075	10	4031	965	1343	6464
Pseudomonas fluorescens/azotoformans	ATGC076	11	3958	1552	975	7120
Pseudomonas chlororaphis/fluorescens/protegens	ATGC078	16	3418	2388	308	8651
Clostridium botulinum/sporogenes	ATGC081	14	3112	488	1477	4604
Burkholderia pseudomallei/mallei/thailandensis	ATGC088	20	3612	2028	357	4552
Burkholderia cenocepacia/cepacia/multivorans	ATGC089	19	3829	2618	501	11276
Sulfolobus islandicus/solfataricus	ATGC093	20	2230	397	792	1835
Serratia marcescens/liquefaciens/plymuthica	ATGC098	19	3505	1235	228	2895
Vibrio parahaemolyticus/alginoliticus/antiquarius	ATGC110	12	3886	587	65	637
Aeromonas hydrophila/veronii/salmonicida	ATGC111	15	2426	1757	2	3062
Shewanella baltica/putrefaciens/oneidensis	ATGC120	14	2929	1259	66	2217
Rhizobium leguminosarum/etli	ATGC123	10	3883	2362	217	8186
Yersinia pestis/pseudotuberculosis/enterocolitica	ATGC127	20	2420	1535	234	2270
Xanthomonas citri/campestris/axonopodis	ATGC134	13	3014	1119	130	1368
Neisseria meningitidis	ATGC137	15	1521	510	1230	8592
Francisella tularensis/philomiragia/noatunensis	ATGC138	20	1062	617	53	954
Acinetobacter baumannii/pittii	ATGC149	20	2874	715	94	818
Ralstonia solanacearum/mannitolilytica/pickettii	ATGC188	10	2963	1587	837	11668
Alteromonas mediterranea/macleodii	ATGC190	14	2889	943	179	1441
Bartonella henselae/bacilliformis/quintana	ATGC201	12	799	535	1	236
Fusobacterium nucleatum/hwasookii	ATGC261	13	1338	782	106	3503
Bacillus pumilus	ATGC290	10	3102	560	3	435
Streptococcus anginosus/constellatus/intermedius	ATGC296	11	1152	647	1	908

TABLE S1: Genome clusters (ATGCs) in the analyzed dataset and model parameters that were inferred under the IGP-CGS assumptions.

Genomes names	ATGC	2-class R^2	2-class R^2_{adj}	2-class Δg_1	3-class R^2	3-class R^2_{adj}	3-class Δg_1
Escherichia coli/Salmonella enterica	ATGC001	0.999957	0.999952	2771	0.999957	0.999946	2771
Klebsiella/Enterobacter aerogenes	ATGC002	0.999893	0.999881	2714	0.999960	0.999949	3010
Bacillus thuringiensis/cereus/anthracis	ATGC014	0.999924	0.999915	2505	0.999924	0.999904	2505
Bacillus subtilis/amyloliquefaciens/velezensis	ATGC015	0.999865	0.999849	1291	0.999865	0.999829	1291
Mycoplasma mycoides/capricolum	ATGC035	0.999635	0.999583	502	0.999635	0.999513	502
Rickettsia rickettsii/conorii/massiliae	ATGC044	0.999808	0.999777	384	0.999808	0.999732	384
Ehrlichia chaffeensis/ruminantium/muris	ATGC045	0.999985	0.999982	58	0.999985	0.999978	58
Helicobacter pylori/cetorum	ATGC050	0.999945	0.999938	691	0.999946	0.999932	696
Corynebacterium glutamicum/deserti/callunae	ATGC069	0.995957	0.995222	962	0.999915	0.999878	1328
Pseudomonas putida/monteilii/parafulva	ATGC071	0.999880	0.999866	3885	0.999941	0.999926	4107
Pseudomonas syringae/savastanoi/cichorii	ATGC075	0.999993	0.999991	494	0.999993	0.999988	494
Pseudomonas fluorescens/azotoformans	ATGC076	0.999885	0.999856	2275	0.999885	0.999808	2275
Pseudomonas chlororaphis/fluorescens/protegens	ATGC078	0.999880	0.999861	2102	0.999880	0.999836	2102
Clostridium botulinum/sporogenes	ATGC081	0.999865	0.999841	1128	0.999865	0.999805	1128
Burkholderia pseudomallei/mallei/thailandensis	ATGC088	0.999674	0.999636	2715	0.999674	0.999587	2715
Burkholderia cenocepacia/cepacia/multivorans	ATGC089	0.999870	0.999854	5515	0.999870	0.999833	5515
Sulfolobus islandicus/solfataricus	ATGC093	0.999943	0.999936	1502	0.999944	0.999929	1522
Serratia marcescens/liquefaciens/plymuthica	ATGC098	0.999707	0.999671	1866	0.999707	0.999624	1866
Vibrio parahaemolyticus/alginolyticus/antiquarius	ATGC110	0.999918	0.999900	1005	0.999964	0.999943	1130
Aeromonas hydrophila/veronii/salmonicida	ATGC111	0.999472	0.999384	1358	0.999925	0.999896	1552
Shewanella baltica/putrefaciens/oneidensis	ATGC120	0.999896	0.999877	2140	0.999896	0.999850	2140
Rhizobium leguminosarum/etli	ATGC123	0.999790	0.999731	2456	0.999790	0.999623	2456
Yersinia pestis/pseudotuberculosis/enterocolitica	ATGC127	0.999859	0.999843	2045	0.999859	0.999822	2045
Xanthomonas citri/campestris/axonopodis	ATGC134	0.999837	0.999804	889	0.999837	0.999755	889
Neisseria meningitidis	ATGC137	0.999752	0.999710	700	0.999752	0.999652	700
Francisella tularensis/philomiragia/noatunensis	ATGC138	0.999873	0.999858	1101	0.999873	0.999840	1109
Acinetobacter baumannii/pittii	ATGC149	0.999874	0.999860	2515	0.999891	0.999862	2577
Ralstonia solanacearum/mannitolilytica/pickettii	ATGC188	0.999997	0.999996	-4	0.999997	0.999994	-4
Alteromonas mediterranea/macleodii	ATGC190	0.999811	0.999776	1857	0.999811	0.999726	1857
Bartonella henselae/bacilliformis/quintana	ATGC201	0.999768	0.999717	113	0.999983	0.999973	151
Fusobacterium nucleatum/hwasookii	ATGC261	0.999712	0.999654	1040	0.999712	0.999568	1040
Bacillus pumilus	ATGC290	0.999601	0.999487	566	0.999798	0.999637	609
Streptococcus anginosus/constellatus/intermedius	ATGC296	0.999738	0.999673	587	0.999738	0.999564	587

TABLE S2: Comparison of 2-class and 3-class model fits. Goodness of fit R^2 , adjusted goodness of fit R^2_{adj} , and the difference between model prediction for the number of singletons and the actual value Δg_1 are listed. Eight cases where the 3-class fit is better than the 2-class fit in terms of R^2_{adj} are highlighted. In all cases, the error in the 3-class model prediction for the number of singletons is greater or equal to the error of the 2-class model prediction.




FIG. S1: Genome intersections and gene commonality distribution for the analyzed genomic dataset. The IGP-CGS model fits are also indicated, see legend of Fig. 4 in the main text. Each row shows a different cluster, which is indicated in the row heading.




FIG. S2: Same as Fig. S1.




FIG. S3: Same as Fig. S1.




FIG. S4: Same as Fig. S1.




FIG. S5: Same as Fig. S1.




FIG. S6: Same as Fig. S1.




FIG. S7: Genome intersections, gene cumulative commonality and gene commonality distribution for the analyzed genomic dataset. The IGP-CGS model fits are indicated, as shown in the upper left panel legend.




FIG. S8: Same as Fig. S7.




FIG. S9: Same as Fig. S7.




FIG. S10: Same as Fig. S7.




FIG. S11: Same as Fig. S7.




FIG. S12: Same as Fig. S7.




FIG. S13: Comparison of the statistics of the IGP-CGS model fit to the 33 genomic clusters, when fitted to mean intersections $\langle I \rangle_k$, gene commonality g_n , and gene cumulative commonality J_k . **a)** Histogram for the error in core-genome sizes g_N of model fit. The error is calculated as $(g_N^{\text{model}} - g_N^{\text{data}})/g_N^{\text{data}}$. **b)** Histogram of the error in model prediction for the number of singletons, as computed from model mean genomes intersections using Eq. 3. The error is calculated as in panel **a**.




FIG. S14: The similarity of simulated datasets to the genomic data of ATGC001. Simulations for different pool sizes under the FGP-CGS assumption are shown in panels **a-d**. Simulations for different gain to loss ratios under the IGP-VGS assumption are shown in panels **e-h**. The similarity between the simulated data is quantified by the error in the number of singletons (panels **a** and **e**), the error in the pangenome size (panels **b** and **f**), a combined measure of the number of singletons and the pangenome size (panels **c** and **g**), and the goodness of fit for the mean intersections (panels **d** and **h**). The error dX is calculated as $(X_{\text{model}} - X_{\text{data}})/X_{\text{data}}$. Contour lines of the optimal combined measure are indicated in panels **c** and **g**. For comparison, the optimal region in terms of the combined measure is also shown in panels **d** and **h**. The parameters that were used in the simulations that are shown in Fig. 8 of the main text are indicated by a vertical and a horizontal black lines in all panels.