

Additional file 3

Spoligotype-Specific Risk Of Finding Lesions In Tissues From Cattle Infected By *Mycobacterium bovis*

Alberto Gómez-Buendía¹, Beatriz Romero¹, Javier Bezos^{1,2}, Francisco Lozano¹, Lucía de Juan^{1,2}, Julio Álvarez^{1,2*}

¹ VISAVET Health Surveillance Centre, Universidad Complutense de Madrid, Spain

² Departamento de Sanidad Animal, Facultad de Veterinaria, Universidad Complutense de Madrid, Spain

*Correspondence: jalvarez@visavet.ucm.es

Frequency of isolates belonging to the most frequent spoligotypes cultured from cattle from lesioned mediastinal lymph nodes from cattle depending on the type of lesion found

Spoligotype	Type of lesion			Total
	1	2	3	
SB0121	83 (32.4%)	61 (23.8%)	112 (43.6%)	256
SB0339	49 (34.8%)	37 (27.4%)	51 (37.8%)	135
SB0134	19 (27.1%)	23 (32.9%)	28 (40.0%)	70
SB0265	19 (30.7%)	18 (29.0%)	25 (40.3%)	62
SB0120	23 (43.4%)	13 (24.5%)	17 (32.1%)	53
SB0295	7 (14.9%)	10 (21.3%)	30 (63.8%)	47
SB1142	9 (29.0%)	9 (29.0%)	13 (42.0%)	31
Total	207 (31.7%)	171 (26.2%)	276 (42.2%)	654

Frequency of isolates belonging to the most frequent spoligotypes cultured from cattle from lesioned lungs from cattle depending on the type of lesion found

Spoligotype	Type of lesion					Total
	1	2	3	4	5	
SB0121	17 (19.3%)	25 (28.4%)	18 (20.5%)	10 (11.4%)	18 (20.5%)	88
SB0339	6 (13.0%)	8 (17.4%)	18 (39.1%)	7 (15.2%)	7 (15.2%)	46
SB0134	2 (6,3%)	3 (9.4%)	8 (25.0%)	9 (28.1%)	10 (31.3%)	32
SB0265	1 (4.6%)	5 (22.7%)	6 (27.3%)	5 (22.7%)	5 (22.7%)	22
SB0120	2 (11.8%)	1 (5.9%)	7 (41.2%)	5 (29.4%)	2 (11.8%)	17
SB0295	0 (0.0%)	2 (20.0%)	3 (30.0%)	1 (10.0%)	4 (40.0%)	10
SB1142	2 (20.0%)	3 (30.0%)	3 (30.0%)	0 (0.0%)	2 (20.0%)	10
Total	30 (13.3%)	47 (20.9%)	63 (28.0%)	36 (16.0%)	49 (21.8%)	225