

Additional file 1

Electronic Cigarette Menthol Flavoring is Associated with Increased Inhaled Micro and Sub-micron Particles and Worse Lung Function in Combustion Cigarette Smokers

AUTHORS

Divay Chandra^{1#}, Rachel F. Bogdanoff^{1#}, Russell P. Bowler², Kambez H. Benam^{1,3,4*}

AFFILIATIONS

¹Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, University of Pittsburgh, Pittsburgh, PA 15213, USA.

²Division of Pulmonary, Critical Care and Sleep Medicine, National Jewish Health, Denver, CO 80206, USA.

³Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA 15219, USA.

⁴Vascular Medicine Institute, University of Pittsburgh, Pittsburgh, PA 15213, USA.

#: These authors contributed equally.

CONTACT

*Corresponding author: benamk@pitt.edu

PG:VG Ratio	50:50 PG:VG (mL)	VG (mL)
38:62	19.0	6.0
41:59	20.5	4.5
47:53	23.5	1.5
48:52	24.0	1.0
50:50	25.0	0

**Table S1. Summary of PG and VG Volumes Used to Create
PG:VG Controls for Vuse E-liquid in Studies in Fig. 2.**