

Supplementary Figure 1 The role of lipids droplets in ferroptosis induced by RSL3 in GC cells. **A** Heatmap of the spatial distribution of PUFAs in various lipid contents of cells cultured with attachment and detachment analyzed by lipid mass spectrometry. **B** BODIPY 493/503 imaging of neutral lipid accumulation (green) in GC cells treated with OA (200 μ M) or co-culture with adipocytes. Cell nuclei were stained with DAPI. Scale bars 10 μ m. **C** Cell viability assessment of indicated GC cells cultured alone or with adipocytes and treated with RSL3 for 24 h. **D** Cell viability assessment of indicated GC cells treated with AA (20 μ M) and OA (200 μ M) for 24 h under attached conditions. Data are mean $\pm$ SD of at least three replicates from two independent experiments. P values were determined by a two-tailed t-test.

Supplementary Figure 2 GSEA analysis of the activation and expression of gene pathways in cancer tissues from patients with high and low GPD1/1L expression (**A, B**).


Supplementary Figure 3 The effect of lipid droplets on FSP1. **A** Western blot analysis of FSP1 and GPX4 expression in GC cells co-cultured with adipocytes for 24 h. **B** Western blot analysis of FSP1, GPD1 and GPD1L expression in GPD1/1L knockout GC cells treated with OA (200 μ M) for 12 h and 24 h.

A




Supplementary Figure 2

A



Rank gene list, GPD1 up verse down


B



Rank gene list, GPD1L up verse down

Supplementary Figure 3

A



B

