

Supplementary Table S1. Fatigue score according to JIA category in the Nordic JIA cohort at 18-year follow-up

JIA category ^a	No. assessed	Fatigue ^b mean $\pm$ SD	Severe fatigue ^c No. (%)
Systemic	13	3.2 $\pm$ 1.6	3 (23)
Oligo. persistent	98	3.0 $\pm$ 1.4	25 (26)
Oligo. extended	77	3.2 $\pm$ 1.6	23 (30)
Polyart. RF negative	64	3.1 $\pm$ 1.3	13 (20)
Polyart. RF positive	6	4.4 $\pm$ 1.4	3 (50)
Psoriatic	21	3.4 $\pm$ 1.6	5 (24)
ERA	37	3.1 $\pm$ 1.3	8 (22)
Undifferentiated	61	3.4 $\pm$ 1.7	19 (31)

JIA = juvenile idiopathic arthritis; No. = numbers; SD = standard deviation; Oligo. = oligoarticular; Polyart. = polyarticular; RF = rheumatoid factor; ERA = enthesitis-related arthritis

^a JIA category according to the International League of Associations for Rheumatology (ILAR) classification criteria

^b Fatigue measured with Fatigue Severity Scale global score, 1-7 (1= lowest, 7 = highest)

^c Fatigue Severity Scale $\geq$ 4