



Uncropped gels for Western Blots in Figure 3

STAT3 : 88 kDa Tubulin: 50 kDa




RORyt: 34 kDa β -actin: 42 kDa




CD4⁺ T Cell lysate, 4-20% SDS-PAGE gel, MOPS running buffer.

Uncropped gels for Western Blots in Figure 4

STAT3: 88 kDa β -actin: 42 kDa




RORyt: 58 kDa GAPDH: 36 kDa




Uncropped gels for Western Blots in Figure 5

Claudin 5: 18kDa GAPDH: 36 kDa




Occludin: 59 kDa GAPDH: 36 kDa




Mouse brain tissue lysate, 4-20% SDS-PAGE gel, MOPS running buffer.

Uncropped gels for Western Blots in Figure 6


p-JNK: 48\53 kDa GAPDH: 36 kDa




JNK: 48\53 kDa GAPDH: 36kDa




p-ERK: 41\43 kDa Tubulin: 50 kDa




ERK: 41\43 kDa Tubulin: 50 kDa



p-P38: 41 kDa Tubulin: 50 kDa




P38: 41 kDa Tubulin: 50 kDa




Mouse brain tissue lysate, 4-20% SDS-PAGE gel, MOPS running buffer.

Uncropped gels for Western Blots in Figure 6

PSD95: 80 kDa Tubulin: 50 kDa




Synapsin-1: 77 kDa Tubulin: 50kDa




Uncropped gels for Western Blots in Figure S3

IL-17RA: 135 kDa Tubulin: 50 kDa



P38: 41 kDa Tubulin: 50 kDa



p-P38: 41 kDa Tubulin: 50 kDa

