

Additional File 1: Copy of the survey questions

What is your age?

26-30

31-35

36-40

41-45

46-50

51-55

56-60

61-65

>65

What school/university/college did you complete your chiropractic education at?

NSW

VIC

WA

New Zealand

Other (please specify) _____

How many years have you been in practice?

6-10

11-15

16-20

21-30

31-40

>40

In which state do you practice? (you may select more than one response if appropriate)

NSW

VIC

SA

ACT

QLD

TAS

WA

NT

New Zealand

Other (Please specify) _____

How often do you apply the following in your practice?

	Always	Most of the Time	Sometimes	Rarely	Never
Wellness care					
Subluxation-based care					
Treatment of musculoskeletal dysfunction					
Treatment of musculoskeletal pain					
Evidence informed practice					
Rehabilitation or exercise prescription					
Other (Please specify)					

Which technique system do you mainly use in your practice?

Diversified

Gonstead

Thompson/TPT

Instrument (Activator)

Other (Please specify) _____

In the following questions: for each of the musculoskeletal conditions listed please select your most commonly used treatment modalities from the drop-boxes provided. If you have more than one, please select them in order of preference. If you have only one/two preferences you can leave the other box(es) blank.

Cervical myofascial pain syndrome

[illegible]

Torticollis

[illegible]

Cervical facet syndrome

[illegible]

Cervical disc syndrome (with radiculopathy)

[illegible]

Cervical disc syndrome (without radiculopathy)

[illegible]

Cervical lateral stenosis

[illegible]

Cervical central stenosis

[illegible]

Cervical related headache

[illegible]

Thoracic myofascial pain syndrome

[illegible]

Thoracic facet syndrome

[illegible]

Rib dysfunction

[illegible]

Lumbar myofascial pain syndrome

[illegible]

Lumbar facet syndrome

[illegible]

Lumbar disc syndrome (without radiculopathy)

[illegible]

Lumbar disc syndrome (with radiculopathy)

[illegible]

Lumbar lateral stenosis

[illegible]

Lumbar central stenosis

[illegible]

Sacroiliac joint dysfunction

	Diversified	Gonstead	Thompson/TPT	Flexion Distraction	Instrument (Activator or similar)	Soft tissue therapy	Electrophysical therapy	Exercise prescription	Other
1st most commonly used treatment modality									
2nd most commonly used treatment modality									
3rd most commonly used treatment modality									

If you selected the response 'other' as a treatment modality in any of the above questions, please use the space below to describe the treatment modality that you would use.