


Study	EBF	Early initiation	STSC
<i>Odu et al. [27]</i>	73.8	75	
<i>Aliyu et al. [19]</i>	70.1		
<i>Yakubu et al. [21]</i>	70		
<i>Mohammed & Aliyu, [26]</i>	68.5		
<i>Elegbua et al. [31]</i>	60.3	58.4	
<i>Akadri & Odelola, [20]</i>	58.8	38.8	
<i>Okoroiwu et al. [28]</i>	54.4	99.3	
<i>Balogun et al. [30]</i>	52.4	59.6	
<i>Olasinde et al. [29]</i>	46.1	40.6	
<i>Anazonwu et al. [34]</i>	39.9		
<i>Anaba et al. [22]</i>	37.5	42.1	29.8
<i>Atimati & Adam, [24]</i>	36.6	44.5	
<i>Ugboaja et al. [35]</i>	35.9		
<i>Adamu et al. [18]</i>	34.2		
<i>Ogundairo et al. [36]</i>	33.1		
<i>Bisi-Onyemaechi et al. [25]</i>	26		
<i>Anyanwu et al. [23]</i>	25		
<i>Amat Camacho et al. [32]</i>	12.5		



Supplementary Table 2 with Figure 3: Summary of Prevalence Rate of EBF, EI

Joseph & Earland, [33]

Low EBF

Low EI



3F
early initiation
TSC



