

Additional Files: Supplementary Figures

Loss of endosomal recycling factor RAB11 coupled with complex regulation of MAPK/ERK/AKT signaling in postmortem spinal cord specimens of sporadic amyotrophic lateral sclerosis patients

Joy Mitra¹, Pavana M. Hegde¹ and Muralidhar L. Hegde^{1,2,3*}

¹Department of Radiation Oncology, Houston Methodist Research Institute, Houston, TX 77030, USA; ²Weill Medical College, New York, NY 10065, USA; ³Houston Methodist Neurological Institute, Institute of Academic Medicine, Houston Methodist, Houston, TX 77030, USA

* Corresponding Author: Muralidhar L. Hegde (mlhegde@houstonmethodist.org)

Email addresses of authors:

Joy Mitra (jmitra@houstonmethodist.org)

Pavana M. Hegde (pdixit@houstonmethodist.org)

Muralidhar L. Hegde (mlhegde@houstonmethodist.org)

Supplementary Material included two Additional Figures.

Additional file 1: Figure S1. Relative levels of monomeric TDP-43 and RAB11 in ALS patients spinal cord tissue. [Related to Fig. 1a.](#) Histogram plot showing relative expression levels of monomeric TDP-43 and RAB11 in spinal cord tissue extracts from four controls and 10 ALS patients. X-axis indicates sample number and Y-axis denotes relative protein level, quantified from wester blots. The TDP-43 monomer levels were previously reported in Mitra et al. [8] and RAB11 levels are shown in Fig. 1a in this study.

