

Appendix 2. Standardized Toxicity Tables

HEMATOLOGY				
	Grade 1 (Mild)	Grade 2 (Moderate)	Grade 3 (Severe)	Grade 4 (Potentially life threatening)
Hemoglobin	9.5 - 10.5 gm/dL	8.0 - 9.4gm/dL	6.5 - 7.9 gm/dL	< 6.5 gm/dL
Absolute Neutrophil Count	1000-1500/mm ³	750-999/mm ³	500-749/mm ³	<500/mm ³
Platelets	75,000-99,999/mm ³	50,000-74,999/mm ³	20,000-49,999/mm ³	<20,000/mm ³
WBCs	11,000-13,000/mm ³	13,000- 15,000 /mm ³	15,000-30,000/mm ³	>30,000 or <1,000 /mm ³
% Polymorphonuclear Leucocytes + Band Cells	> 80%	90 – 95%	>95%	-----
CHEMISTRIES				
	Grade 1	Grade 2	Grade 3	Grade 4
Hyponatremia	130 to<135 mEq/L	123-129 mEq/L	116-122 mEq/L	< 116 mEq/L or abnormal sodium <i>with</i> mental status changes or seizures
Hypernatremia	146-150 mEq/L	151-157 mEq/L	158-165 mEq/L	> 165 mEq/L or abnormal sodium <i>with</i> mental status changes or seizures
Hypokalemia	3.0 - 3.4 mEq/L	2.5 - 2.9 mEq/L	2.0 - 2.4 mEq/L or intensive replacement therapy or hospitalization required	< 2.0 mEq/L or abnormal potassium <i>with</i> paresis, ileus, or life-threatening arrhythmia
Hyperkalemia	5.6 - 6.0 mEq/L	6.1 - 6.5 mEq/L	6.6 - 7.0 mEq/l	> 7.0 mEq/L or abnormal potassium <i>with</i> life-threatening arrhythmia
Hypoglycemia	55-64 mg/dL	40-54 mg/dL	30-39 mg/dL	<30 mg/dL or abnormal glucose <i>with</i> mental status changes or coma
Hyperglycemia (non-fasting and no prior diabetes)	116 - 160 mg/dL	161- 250 mg/dL	251 - 500 mg/dL	> 500 mg/dL or abnormal glucose <i>with</i> ketoacidosis or seizures
Creatinine	1.1 - 1.5 x ULN	1.6 - 3.0 x ULN	3.1 - 6 x ULN	> 6 x ULN or dialysis required
ENZYMES				
	Grade 1	Grade 2	Grade 3	Grade 4
AST (SGOT)	1.1 - <2.0 x ULN	2.0 – <3.0 x ULN	3.0 – 8.0 x ULN	> 8 x ULN
ALT (SGPT)	1.1 - <2.0 x ULN	2.0 – <3.0 x ULN	3.0 – 8.0 x ULN	> 8 x ULN
URINALYSIS				
	Grade 1	Grade 2	Grade 3	Grade 4
Proteinuria	1+ or 200 mg - 1 gm loss/day	2-3+ or 1- 2 gm loss/day	4+ or 2-3.5 gm loss/day	nephrotic syndrome or > 3.5 gm loss/day

Hematuria	microscopic only <10 rbc/hpf	gross, no clots >10 rbc/hpf	gross, with or without clots, OR red blood cell casts	obstructive or required transfusion
CARDIOVASCULAR				
	Grade 1	Grade 2	Grade 3	Grade 4
Cardiac Rhythm	asymptomatic, no Rx required	asymptomatic, transient signs, no Rx required	recurrent/persiste nt; symptomatic Rx required	unstable dysrhythmia; hospitalization and treatment required
Hypertension	transient increase > 20 mm/Hg; no treatment	recurrent, chronic increase > 20mm/Hg. /treatment required	acute treatment required; outpatient treatment or hospitalization possible	end organ damage or hospitalization required
Hypotension	transient orthostatic hypotension with heart rate increased by <20 beat/min or decreased by <10 mm Hg systolic BP, No treatment required	symptoms due to orthostatic hypotension or BP decreased by <20 mm Hg systolic; correctable with oral fluid treatment	requires IV fluids; no hospitalization required	mean arterial pressure <60mm/ Hg or end organ damage or shock; requires hospitalization and vasopressor treatment
Pericarditis	minimal effusion	mild/moderate asymptomatic effusion, no treatment	symptomatic effusion; pain; EKG changes	tamponade; pericardiocentesis or surgery required
Hemorrhage, Blood Loss	microscopic/occ ult	mild, no transfusion	gross blood loss; 1-2 units transfused	massive blood loss; > 3 units transfused
RESPIRATORY				
	Grade 1	Grade 2	Grade 3	Grade 4
Cough	transient- no treatment	persistent cough; treatment responsive	Paroxysmal cough; uncontrolled with treatment	-----
Bronchospasm, Acute	transient; no treatment; 70% - 80% FEV1 of peak flow	requires treatment; normalizes with bronchodilator; FEV1 50% - 70% (of peak flow)	no normalization with bronchodilator; FEV1 25% - 50% of peak flow; or retractions present	cyanosis: FEV1 < 25% of peak flow or intubation necessary
Dyspnea	dyspnea on exertion	dyspnea with normal activity	dyspnea at rest	dyspnea requiring hospitalization
GASTROINTESTINAL				
	Grade 1	Grade 2	Grade 3	Grade 4
Nausea	mild or transient; maintains reasonable intake	moderate discomfort; intake decreased significantly; some activity limited	no significant intake; requires IV fluids	hospitalization required;

Vomiting	1 episode in 24 hours	2-5 episodes in 24 hours	>6 episodes in 24 hours or needing IV fluids	physiologic consequences requiring hospitalization or requiring parenteral nutrition
Constipation	requiring stool softener or dietary modification	requiring laxatives	constipation requiring manual evacuation or enema	obstruction or toxic megacolon
Diarrhea	mild or transient; 3-4 loose stools/day or mild diarrhea last < 1 week	moderate or persistent; 5-7 loose stools/day or diarrhea lasting >1 week	>7 loose stools/day or bloody diarrhea; or orthostatic hypotension or electrolyte imbalance or >2L IV fluids required	hypotensive shock or physiologic consequences requiring hospitalization
Oral Discomfort/Dysphagia	mild discomfort; no difficulty swallowing	some limits on eating/drinking	eating/talking very limited; unable to swallow solid foods	unable to drink fluids; requires IV fluids
NEUROLOGICAL				
	Grade 1	Grade 2	Grade 3	Grade 4
Neuro-Cerebellar	slight incoordination dysdiadochokinesis	intention tremor, dysmetria, slurred speech; nystagmus	locomotor ataxia	incapacitated
Psychiatric	mild anxiety or depression	moderate anxiety or depression; therapy required; change in normal routine	severe mood changes requiring therapy; or suicidal ideation; or aggressive ideation	acute psychosis requiring hospitalization; or suicidal gesture/attempt or hallucinations
Muscle Strength	subjective weakness no objective symptoms/signs	mild objective signs/symptoms, no decrease in function	objective weakness function limited	paralysis
Paresthesia (burning, tingling, etc.)	mild discomfort; no treatment required	moderate discomfort; non-narcotic analgesia required	severe discomfort; or narcotic analgesia required with symptomatic improvement	incapacitating; or not responsive to narcotic analgesia
Neuro-sensory	mild impairment in sensation (decreased sensation, e.g., vibratory, pinprick, hot/cold in great toes) in focal area or symmetrical	moderate impairment (mod decreased sensation, e.g., vibratory, pinprick, hot/cold to ankles) and/or joint position or mild impairment that is not symmetrical	severe impairment (decreased or loss of sensation to knees or wrists) or loss of sensation of at least mod degree in multiple different body	sensory loss involves limbs and trunk; paralysis; or seizures

	distribution; or change in taste, smell, vision and/or hearing		areas (i.e., upper and lower extremities)	
SKIN				
	Grade 1	Grade 2	Grade 3	Grade 4
Mucocutaneous	erythema; pruritus	diffuse, maculopapular rash, dry desquamation	vesiculation or moist desquamation or ulceration	exfoliative dermatitis, mucous membrane involvement or erythema, multiforme or suspected Stevens-Johnson or necrosis requiring surgery
Induration	< 15mm	15-30 mm	>30mm	
Erythema	< 15mm	15-30 mm	>30mm	
Edema	< 15mm	15-30 mm	>30mm	
Rash at Injection Site	< 15mm	15-30 mm	>30mm	
Pruritus	slight itching at injection site	moderate itching at injection extremity	itching	
SYSTEMIC				
	Grade 1	Grade 2	Grade 3	Grade 4
Allergic Reaction	pruritus without rash	localized urticaria	generalized urticaria; angioedema	anaphylaxis
Headache	mild, no treatment required	transient, moderate; treatment required	Severe; responds to initial narcotic therapy	intractable; requires repeated narcotic therapy
Fever: oral	37.7 - 38.5 C or 100.0 - 101.5 F	38.6 - 39.5 C or 101.6 - 102.9 F	39.6 - 40.5 C or 103 - 105 F	> 40 C or > 105 F
Fatigue	normal activity reduced < 48 hours	normal activity decreased 25- 50% > 48 hours	normal activity decreased > 50% can't work	unable to care for self

Additionally, table below* describes a severity grading system for any clinical adverse event not identified in appendix 4, standardized toxicity table.

Grade 1 (mild)	Grade 2 (moderate)	Grade 3 (severe)	Grade 4 (potentially life-threatening)
Mild symptoms causing no or minimal interference with usual social & functional activities with intervention not indicated	Moderate symptoms causing greater than minimal interference with usual social & functional activities with intervention indicated	Severe symptoms causing inability to perform usual social & functional activities with intervention or hospitalization indicated	Potentially life- threatening symptoms causing inability to perform basic self-care functions with intervention indicated to prevent permanent impairment, persistent disability, or death

*Adapted from the NIH Division of AIDS (DAIDS) Table for Grading the Severity of Adult and Pediatric Adverse Events.