

Table S1. Effect of HEWs led home-based multicomponent intervention on linkage to hypertension care, northwest Ethiopia, March 2021-January 2022: Subgroup analysis

Characteristics	Stratum specific RBI (95% CI)	P value
Sex		
Male	1.77 (1.38, 2.26)	<0.00
Female	1.60 (1.27, 2.01)	<0.001
Age, years		
25-34	3.43 (1.16, 10.19)	0.026
35-44	2.00 (1.32, 3.03)	0.001
45-54	1.35 (1.00, 1.81)	0.047
55-64	1.69 (1.14, 2.52)	0.009
≥65	1.65 (1.24, 2.20)	0.001
Educational status		
Could not read or write	1.82 (1.50, 2.22)	<0.00
Able to read and write	1.10 (0.77, 1.57)	0.589
Primary school and above	1.95 (0.90, 4.20)	0.089
Stage of hypertension		
Stage 1	2.32 (1.45, 3.72)	<0.001
Stage 2	1.56 (1.30, 1.87)	<0.001
Family history of hypertension		
Yes	1.15 (0.84, 1.58)	0.388
No	1.86 (1.52, 2.26)	<0.001
Community-based health insurance		
Adequately insured	1.54 (1.15, 2.06)	0.004
Partially insured	1.02 (0.66, 1.59)	0.913
Not insured	1.87 (1.49, 2.35)	<0.001