

Table S5. Host blood meal identification of *Culicoides* biting midges collected from Ron Phibun district in Nakhon Si Thammarat, southern Thailand

Species	Number of blood-fed females	Single blood meal (<i>n</i>)					Multiple blood meals (<i>n</i>)
		Human	Cow	Dog	Pig	Chicken	
<i>C. actoni</i>	2	0	2	0	0	0	-
<i>C. fulvus</i>	3	0	3	0	0	0	-
<i>C. huffi</i>	1	0	0	0	0	1	-
<i>C. innoxius</i>	25	0	24	0	0	0	Cow and chicken (1)
<i>C. insignipennis</i>	18	0	18	0	0	0	-
<i>C. jacobsoni</i>	14	0	14	0	0	0	-
<i>C. orientalis</i>	12	0	12	0	0	0	Human and cow (1)
<i>C. oxystoma</i>	5	0	4	0	0	0	Human and cow (1)
<i>C. peregrinus</i>	10	0	8	0	0	0	-
<i>C. shortti</i>	1	0	1	0	0	0	-
<i>C. sumatrae</i>	12	0	11	0	0	0	Human and cow (1)
<i>C. subgenus Trithecoides</i>	56	0	53	3	0	0	-
Total	159	0	149	3	0	1	4