

Additional file 1

Table S1. Proportion of flies carrying helminth eggs, summarized by species, location, and sex.

Location	Sex	Species	
		<i>Ascaris suum</i> , n/N (%)	<i>Taenia saginata</i> , n/N (%)
Exoskeleton	Female	1/21 (4.8%)	2/21 (9.5%)
	Male	1/27 (3.7%)	1/27 (3.7%)
	Total	2/48 (4.2%)	3/48 (6.3%)
Digestive tract	Female	13/21 (61.9%)	14/21 (66.7%)
	Male	5/27 (18.5%)	17/27 (63.0%)
	Total	18/48 (37.5%)	31/48 (64.6%)

Abbreviations: n, number of positive flies; N, total number of flies examined.

Table S2. Total and median egg counts recovered from flies, summarized by species, location, and sex.

Location	Sex	Species					
		<i>Ascaris suum</i>			<i>Taenia saginata</i>		
		Total count (min–max) ^a	Median count, all flies (Q1–Q3) ^b	Median count, positive flies (Q1–Q3) ^c	Total count (min–max) ^a	Median count, all flies (Q1–Q3) ^b	Median count, positive flies (Q1–Q3) ^c
Exoskeleton	Female	1 (1–1)	0 (0–0)	1 (1–1)	2 (1–1)	0 (0–0)	1 (1–1)
	Male	1 (1–1)	0 (0–0)	1 (1–1)	1 (1–1)	0 (0–0)	1 (1–1)
	Total	2 (1–1)	0 (0–0)	1 (1–1)	3 (1–1)	0 (0–0)	1 (1–1)
Digestive tract	Female	35 (1–9)	1 (0–2)	2 (2–3)	103 (1–19)	3 (0–9)	6 (3–10.8)
	Male	9 (1–2)	0 (0–0)	2 (2–2)	75 (1–10)	1 (0–4.5)	4 (1–7)
	Total	44 (1–9)	0 (0–2)	2 (2–2)	178 (1–19)	3 (0–6.2)	5 (3–8)

^a Total egg count recovered from all flies within each subgroup; range indicates minimum–maximum number of eggs per individual fly.

^b Median egg count for all flies; Q1–Q3 represent the first and third quartiles (interquartile range).

^c Median egg count for positive flies; Q1–Q3 represent the first and third quartiles (interquartile range).