

Supporting Information

hAECs Restore Follicular Development in Premature Ovarian Insufficiency via IGFBP2/IGF1R-Mediated Intercellular Communication

Wenjiao Cao^{1#}, Lu Shen^{2#}, Qinyu Zhang¹, Yating Huang¹, Junyan Sun¹, Zixin Cheng¹,
Jing Xu³, Qiuwan Zhang^{1*}, Dongmei Lai^{1*}

¹ International Peace Maternity and Child Health Hospital, Shanghai Key Laboratory of Embryo Original Diseases, School of Medicine, Shanghai Jiao Tong University, Shanghai, 200030, China

² Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders, Ministry of Education, Shanghai Jiao Tong University, Shanghai, 200030, China.

³ School of Biomedical Engineering, Shanghai Jiao Tong University, Shanghai, 200030, China

*Corresponding Author:

Dongmei Lai, MD, PhD

E-mail: laidongmei@hotmail.com

Qiuwan Zhang, PhD

E-mail: zhangqiuwan@163.com

145, Guang-Yuan Road, Shanghai 200030, P. R. China,

Tel: 86-21-64070434

Fax: +86-21-64074642

The original, uncropped Western blot images corresponding to Figure 6 are shown below. Protein molecular weight markers (kDa) are indicated. Areas within red boxes represent the cropped regions used in the Figure 6.

Figure 6A:




Figure 6D:




Figure 6F:

