

Table S1: Primers used for RT-qPCR analysis

Gene symbol	Forward primer	Reverse primer
<i>Arntl/Bmal1</i>	CTATTACCAGGGCAGGCTCA	TTGCTGCCTCATCGTACTG
<i>Clock</i>	TCCATCCAGTATGCCACAGA	CCCACAAGCTACAGGAGCAG
<i>Cry1</i>	GTGGGAAGCGTCCTAGTCAG	ACCAACTTCAGTTGCGAGGA
<i>Cry2</i>	CCATCGTCAATCATGCAGAG	CCACACAGGAAGGGACAGAT
<i>Dbp</i>	AGGAACTGAAGCCTCAACCA	TGGCTGCTTCATTGTTCTTG
<i>Fhl2</i>	CCATTGGAACCAAGAGCTTC	TGCTCCCGGTAAGTAACACC
<i>Hlf</i>	GCTCGCAAAGTCTTCATTCC	CGATCTGGTTCCTCTCAGC
<i>Nfil3/(E4bp4)</i>	GTTACAGCCGCCCTTTCTTT	CATCCATCAATGGGTCCTTC
<i>Npas2</i>	GCCCTCAGAAGTCAGCAGAA	TGCTGTTGGTAGGGTGTGAG
<i>Nr1d1</i>	CTTCCGTGACCTTTCTCAGC	TCACTGTCTGGTCCTTCACG
<i>Nr1d2</i>	CTGGCTTCCGAGATCTGTCT	ACAGTCCGTTCTTTGCATC
<i>Per1</i>	CTGGGGACCAGGTCATTAAG	CTTCAGCACAGAGGCTGCAT
<i>Per2</i>	TTCGGCACATCCCAAAGT	CTCCGTGTCTGGGATCATTT
<i>Per3</i>	AGAAGCTCCAGAGCATGGAA	TCTGTCTTCACAGGCGACAC
<i>Pln</i>	CACTGTGACGATCACCGAAG	TTTCCATTATGCCAGGAAGG
<i>Rora</i>	GAACCACCGAGAAGATGGAA	GGAAAATGGAGTCGCACAAT
<i>Rps16</i>	GCTACCAGGGCCTTTGAGATG	AGGAGCGATTTGCTGGTGTGG
<i>Tef</i>	GTTTGCAGAGGAGGACCTGA	GAGCGTTTAGCTGCCACATT
<i>Ttn</i>	CACACAGCCCAGATCTCAAA	TTCTGGAGCGACTCACACTG