

Table S1. Antimicrobial susceptibility and MIC results of *K. pneumoniae* k9. Antimicrobial susceptibility was performed using the disk diffusion technique; MICs of ertapenem, imipenem and meropenem were determined using the E-test methodology; MIC of colistin was determined using the broth microdilution method. Antibiotics were divided by their drug categories. GM: gentamycin 10ug; IP: imipenem; MEM: meropenem 10ug; IPM: imipenem 10ug; PM: ceftazidime; CTX: cefotaxime 30ug; CXM: cefuroxime 30ug; FEP: cefepime 30ug; CZD: ceftazidime 10ug; CZN: ceftazolin 30ug; SCF: ceftazidime/sulbactam 75/30ug; FOX: ceftazidime 30ug; NOR: norfloxacin 5ug; CIP: ciprofloxacin 5ug; FAD: fusidic acid 10ug; TGC: tigecycline 15ug; CMN: clindamycin 2ug; ERY: erythromycin 15ug; ATM: aztreonam 30ug; LZD: linezolid 30ug; AMX: amoxicillin 25ug; PIR: piperacillin 100ug; AMP: ampicillin 10ug; TZP: piperacillin/tazobactam 100/10ug; FOS: fosfomycin 200ug; TE: tetracycline 30ug; TS: trimethoprim/sulfamethoxazole; CO: colistin; Dark blue: resistant, light blue: intermediate resistance, white: susceptible; MICs are in ug/mL; NA: not available.

[illegible]