

Polymorphisms in DNA repair and oxidative stress genes associated with pre-treatment cognitive function in breast cancer survivors: an exploratory study

Breast Cancer Research and Treatment

Theresa A. Koleck¹ (tat30@pitt.edu), Catherine M. Bender¹, Susan M. Sereika^{1,2}, Adam M. Brufsky³⁻⁵, Barry C. Lembersky^{3,4}, Priscilla F. McAuliffe^{5,6}, Shannon L. Puhalla^{3,5}, Priya Rastogi^{3,5} & Yvette P. Conley^{1,7}

¹School of Nursing, University of Pittsburgh, 3500 Victoria Street, Pittsburgh, PA 15261, USA

²Department of Biostatistics and Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, 130 De Soto Street, Pittsburgh, PA 15261, USA

³Division of Hematology/Oncology, Magee-Womens Hospital of University of Pittsburgh Medical Center (UPMC), 300 Halket Street, Pittsburgh, PA 15213, USA


⁴University of Pittsburgh Cancer Institute, 5150 Centre Avenue, Pittsburgh, PA 15232, USA


⁵School of Medicine, University of Pittsburgh, 3550 Terrace Street, Pittsburgh, PA 15261, USA

⁶Magee-Womens Surgical Associates, Magee-Womens Hospital of University of Pittsburgh Medical Center (UPMC), 300 Halket Street, Pittsburgh, PA 15213, USA

⁷Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, 130 De Soto Street, Pittsburgh, PA 15261, USA

Network 1





© 2000-2015 QIAGEN. All rights reserved.

Online Resource 2 Oxidative stress and DNA repair gene-gene networks generated by pathway analysis

Note. The networks were generated through the use of QIAGEN's Ingenuity Pathway Analysis (IPA[®], QIAGEN Redwood City, www.qiagen.com/ingenuity). The main associated functions of each network are: (Network 1) free radical scavenging, small molecule biochemistry, and neurological disease and (Network 2) DNA replication, recombination, and repair, energy production, and nucleic acid metabolism.