

		<i>Follow up sample</i>	<i>n (%)</i>	<i>Uncontactable or declined</i>	<i>n (%)</i>	<i>Test Statistic</i>
Number in sample			149		208	
Gender	Female		143 (96.0%)		189 (90.8%)	1, 357) = 3.78, p = .06
	Male		6 (4.0%)		19 (9.1%)	
Ethnicity	White British		126 (84.6%)		151 (72.6%)	χ^2 (1, 357) = 7.15, p = .007
	BAME		23 (15.4%)		57 (27.4%)	
Diagnosis	AN		68 (45.6%)		77 (37.0%)	χ^2 (1, 357) = .313, p = .57 ^b
	Atypical AN		51 (34.2%)		94 (45.2%)	
	BN		19 (12.8%)		23 (11.1%)	
	Atypical BN		11 (7.4%)		14 (6.7%)	
Treatment enhanced ^a	OUT		87 (73.1%)		126 (73.7%)	χ^2 (1, 290) = 0.01, p = .91
	AIM		32 (26.9%)		45 (26.3%)	
	Psychiatric admission		5 (3.4%)		7 (3.4%)	
	ITP		19 (12.8%)		24 (11.2%)	
	Admission and ITP		9 (6%)		17 (8.25%)	

		<i>n</i>	<i>Mean (SD)</i>	<i>95% C.I.</i>	<i>n</i>	<i>Mean (SD)</i>	<i>95% C.I.</i>	
Age	At discharge	149	16 yrs 5.3 mths (1 yr, 7.3 mths)	16yrs 1.9 mths - 16yrs 8.3 mths	208	16yrs 1.5mths (1yr, 10.8 mths)	15yrs 10.4mths - 16yrs 4.5mths	F (1, 355) = 2.64, p = .10 d = 0.18
	At follow up	149	23 yrs 3.1 mths (2 yrs 2.9 mths)	22yrs 10.8 mths - 23yrs 7.5mths	201	22yrs 11.8 mths (2yrs 1.8 mths)	22 yrs 8.2 mths – 23yrs 3.2mths	F (1, 348) = 1.36, p = .24, d = -0.13
MCCAED measures	EDE-Q (global) assessment	132	3.54 (1.64)	3.25 – 3.82	173	3.13 (1.73)	2.87 – 3.39	F (1, 303) = 4.34, p = .04, d = -0.24
	EDE-Q (global) discharge	59	1.81 (1.61)	1.39 – 2.23	84	1.49 (1.57)	1.15 – 1.83	F (1, 141) = 1.39, p = .24, d = -0.20
	MFQ assessment	129	33.75 (16.47)	30.88 – 36.61	173	28.37 (15.73)	26.02 – 30.74	F (1, 300) = 8.27, p = .004 d = -0.34
	MFQ discharge	57	16.61 (16.03)	12.36 – 20.87	84	18.59 (16.11)	15.10 – 22.09	F (1, 139) = 0.51, p = .47, d = 0.12

MCCAED biometrics	SCARED assessment	172	28.77 (19.68)	25.38 – 32.16	132	28.29 (17.11)	25.72 – 30.87	F, (1, 302) = 0.05, $p = .82$, $d = -0.03$
	SCARED discharge	59	22.20 (17.39)	17.67 – 26.73	82	21.09 (15.71)	17.63 – 24.54	F, (1, 139) = 0.16, $p = .69$, $d = -0.07$
	%mBMI assessment	147	85.90 (12.25)	83.90 – 87.89	205	86.72 (15.05)	84.65 – 88.79	F, (1, 350) = 0.30, $p = .59$, $d = 0.06$
	%mBMI 3 months	130	88.85 (10.48)	87.04 – 90.67	176	89.35 (11.48)	87.64 – 91.05	F, (1, 304) = 0.15, $p = .70$, $d = 0.05$
	%mBMI discharge	143	93.86 (11.22)	92.00 – 95.71	201	94.63 (13.84)	92.70 – 96.55	F,(1, 342) = 0.30, $p = 0.59$, $d = 0.06$

Supplementary Table S1 | Comparison of those who consented to follow-up and those who were not available to follow-up

^aAN/Atypical AN cases only

^bComparison between AN/Atypical AN and BN/Atypical BN groups

Abbreviations: %mBMI, percentage of median body mass index AIM, Additional Intensive Management (day treatment or inpatient treatment; AN, anorexia nervosa; BAME, Black, Asian or Minority Ethnic; BN, bulimia nervosa; EDE-Q, Eating Disorder Examination Questionnaire global score; MFQ, Mood and Feelings Questionnaire; OPT, Outpatient treatment only; SCARED, Screen for Child Anxiety Related Disorder; SD, standard deviation