

Supplementary Table S10. Sequences of primers used for RT-PCR analyses.

	Forward	Reverse
<i>Ki-67</i>	5'- GACAGAGGTTCTTAAGAGAG -3'	5'- AACAAATCAGATTTGCTTCCG -3'
<i>PCNA</i>	5'- CTGTGTAGTAAAGATGCCTTC -3'	5'- TCTCTATGGTAACAGCTTCC -3'
<i>CCND1</i>	5'- GCCTCTAAGATGAAGGAGAC -3'	5'- CCATTTGCAGCAGCTC -3'
<i>CCNA2</i>	5'- AGTATCATGGTGTTTCTCTCC -3'	5'- AATTTCTACTTGGCCACAAC -3'
<i>ITGB2</i>	5'- TTGAGAAGGAGAAGCTCAAG -3'	5'- CTAACCTCTCAGCAAACCTTGG -3'
<i>ICAM1</i>	5'- GGGGAACCAGAGCCAGGAGACACT -3'	5'- TGGGCCTCACACTTCACTGTCACC -3'
<i>SELL</i>	5'- CAAGAGAAGTATGAATGACCC -3'	5'- TCAGGTAGAAATCTTCCCAG -3'

Ki-67 (Marker of Proliferation Ki-67); *PCNA* (Proliferating Cell Nuclear Antigen); *CCND1* (Cyclin D1); *CCNA2* (Cyclin A2); *ITGB2* (Integrin Subunit Beta 2); *ICAM1* (Intercellular Adhesion Molecule 1); *SELL* (Selectin L)