

Supplementary data, Table 1: Association of clinical factors with overall survival.

BMI, body mass index in kg/m²; ASA, Grading in American Society of Anesthesiologists;

CA 19-9, carbohydrate antigen 19-9

	Overall survival	Lower.CI	Upper.CI	p-value
Age				0.8
≤ 75 years	20.5	17.9	23.0	
> 75 years	20.0	6.8	33.5	
Gender				0.5
Male	23.4	18.3	28.5	
Female	19.1	15.9	22.2	
BMI				0.5
≤ 25 kg/m ²	20.0	17.1	22.9	
> 25 kg/m ²	21.2	16.1	26.3	
CA 19-9				0.2
≤500	20.8	17.8	23.8	
> 500	15.1	11.4	18.8	
Time interval				0.024
< 4 weeks	16.1	12.2	20.1	
≥ 4 weeks	22.9	19.1	26.7	
Adjuvant chemotherapy (n=140)				<0.001
no	9.7	6.4	13.0	
yes	23.8	19.0	29.0	
Tumour stage				0.031
0 + I	23.8	20.0	27.6	
II	20.0	16.8	23.2	
III	17.0	13.9	20.0	
Lymphnode stage				0.1
0	23.7	19.6	27.7	
1	20.8	17.7	24.0	
2	17.0	14.1	19.8	