

Genome Editing for Scalable Production of Alloantigen-free Lentiviral Vectors for *in Vivo* Gene Therapy

Alloantigen-free Lentiviral Vectors

Michela Milani^{1,2}, Andrea Annoni¹, Sara Bartolaccini¹, Mauro Biffi¹, Fabio Russo¹, Tiziano Di Tomaso¹, Andrea Raimondi³, Johannes Lengler⁴, Michael C. Holmes⁵, Friedrich Scheifflinger⁴, Angelo Lombardo^{1,2}, Alessio Cantore^{1*}, Luigi Naldini^{1,2*}

¹San Raffaele Telethon Institute for Gene Therapy, IRCCS San Raffaele Scientific Institute, Milan 20132, Italy;

²Vita Salute San Raffaele University, Milan 20132, Italy;

³IRCCS San Raffaele Scientific Institute, Milan 20132, Italy;

⁴Baxalta (former Baxter) Innovation GmbH, Vienna 1221, Austria;

⁵Sangamo Therapeutics, Inc, Richmond CA 94804, USA;

*These authors share senior authorship


Corresponding author: Luigi Naldini, San Raffaele Telethon Institute for Gene Therapy, Via Olgettina 58, 20132, Milan, Italy; luigi.naldini@hsr.it

Table of Contents


Appendix Figure S1

Appendix Figure S1


A



B



C



Appendix Figure S1 - Generation of LV packaging cell line.

A Western blot on protein extracts from single-cell clones obtained from 293 T-REx stably transfected with pY-Rev, induced (+) or not (-) with dox. Red borders show images taken from different blots.

B LV particles production (ng p24/mL) in the medium of single-cell clones obtained from 293 T-Rex Rev stably transfected with pY-Gag/Pol induced (grey bars) or not (black bars) with dox.

C Western blot on protein extracts from single-cell clones obtained from semi-packaging cell line stably transfected with pY-VSV.G, induced (+) or not (-) with dox. Red borders show images taken from different blots or gels.